February 27, 2023 PEDIATRIC & ADOLESCENT MEDICAL ASSOC PC 117 W Liberty St Rome, NY 13440-5758

Phone: 315-339-0401

Fever Therapy

2-4 MONTHS OF AGE

If your child has a fever, there are several things you can do to bring the temperature down at home. Generally, fevers less than 102° do not require treatment and frequently accompany mild illnesses such as colds, diarrhea, etc. If your child appears unusually ill or if the fever persists, you should consult your doctor. Infants (under 2 months of age) with a rectal temperature of 100° or above should be seen by their physician.

SKIN EXPOSURE: Children lose heat through their skin. Dress the child lightly. If you use sleepers, blankets or

heavy clothes, this will prevent the loss of heat and the fever will go higher.

FLUIDS: Fevers are higher if a child does not have enough fluids in his/her body. Give plenty of

fluids to prevent dehydration and to make the child more comfortable.

ACETAMINOPHEN DOSING--EVERY 4 HOURS

			Infant's Oral	Children's Oral		
		Suspension	Suspension	Chewable	Junior	Adult <u>WEIGHT</u>
AGE	160mg/5mL	160mg/5mL	80mg/tab	160mg/tab	325mg/tab	
6 - 11 LBS	0 - 3 MOS	1.25 mL	_	_		
12 - 17 LBS	4 - 11 MOS	2.5 mL				
18 - 23 LBS	12 - 23 MOS	3.75 mL				
24 - 35 LBS	2 - 3 YRS	5 mL	5mL (1 tsp)	2 tab	1 tab	
36 - 47 LBS	4 - 5 YRS		7.5mL (1.5 tsp)	3 tab	1.5 tab	
48 - 59 LBS	6 - 8 YRS		10mL (2 tsp)	4 tab	2 tab	1 tab
60 - 71 LBS	9 - 10 YRS		12.5mL (2.5 tsp	o) 5 tab	2.5 tab	1 1/4 tab
72 - 95 LBS	11 - 12 YRS		15mL (3 tsp)	6 tab	3 tab	1 1/2 tab

SPONGING:

If the temperature will not return to <103 with the above measures, you may use a lukewarm bath or sponge to lower the fever. *DO NOT USE COLD WATER OR ALCOHOL*. Rub the skin briskly with a washcloth so the blood will come to the surface where it can be cooled. If the child develops chills, stop sponging and in a short while start again. Be sure the water is lukewarm. Bathe the child no more frequently than every 2 hours. It usually take 20-30 minutes in the tub to bring the fever down.